



TURF FERTILIZER

with 0.20% BARRICADE® herbicide

19-0-5

.20% Barricade, PCSCU

**GUARANTEED ANALYSIS**

Total Nitrogen (N) ..... 19.0%

19.0 % Urea nitrogen\*

Soluble Potash (K<sub>2</sub>O) ..... 5.0%

Plant foods derived from: Urea, Poly Coated Sulfur Coated Urea and Muriate of Potash.

\*4.75% slowly available urea nitrogen from poly coated sulfur coated urea.

544-5390

award20sm-14

For selective preemergence control of grass and broadleaf weeds in:

- Established turfgrasses (excluding golf course putting greens), lawns, athletic fields, parks and sod farms
- Field-grown and landscape ornamentals
- Established perennials and wildflower plantings
- Non-crop areas, including landscape beds, ornamentals, and established perennial wildflower plantings on or surrounding municipal grounds and government installations, managed rights-of-way for transportation systems and utilities (including, roadways, roadsides, railways and equipment yards)

**Active Ingredient:**  
 Prodiamine\*.....0.20%  
**Other Ingredients:** .....99.80%  
**Total:** .....100.00%

\*CAS No. 29091-21-2  
 EPA Reg. No. 100-1337-57131  
 EPA Est. No. 57131-WI-1

**KEEP OUT OF REACH OF CHILDREN CAUTION**

See additional precautionary statements and directions for use.

**FIRST AID**

<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

**HOT LINE NUMBER**

For 24 Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), Call 1-800-424-9300.

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals CAUTION**

Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wear protective eyewear. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco. Wear long-sleeved shirt and long pants, shoes, socks, and gloves. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

**Personal Protective Equipment (PPE)**

**Mixers, loaders, applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
  - Shoes plus socks
  - Chemical-resistant gloves made of any waterproof material
- Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Control Statements**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(4-6)), the handler PPE requirements may be reduced or modified as specified in the WPS.

**User Safety Recommendations**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards**

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment wash waters.

**CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, presence of other materials or other influencing factors in the use of the product, which are beyond the control of Eau Claire Coop Oil Co. or Seller. To the extent permitted by applicable law, Buyer and User agree to hold Eau Claire Coop Oil Co. and Seller harmless for any claims relating to such factors.

Eau Claire Coop Oil Co. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. To the extent permitted by applicable law: (1) this warranty does not extend to the use of this product contrary to label instructions or under conditions not reasonably foreseeable to or beyond the control of Seller or Eau Claire Coop Oil Co., and (2) Buyer and User assume the risk of any such use. To the extent permitted by applicable law, Eau Claire Coop Oil Co. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS WARRANTED BY THIS LABEL.

To the extent permitted by applicable law, in no event shall Eau Claire Coop Oil Co. be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **TO THE EXTENT PERMITTED BY APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF Eau Claire Coop Oil Co. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF Eau Claire Coop Oil Co. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

Eau Claire Coop Oil Co. and Seller offer this product, and Buyer and User accept it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of Eau Claire Coop Oil Co..

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirement specific to your State and Tribe, consult the State/Tribal agency responsible for pesticide regulation.

- For use on plants intended for aesthetic purposes or climatic modification.
- Read entire label before using this product.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.** Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**Do not enter or allow others to enter until dusts have settled.**

AWARD®TURF FERTILIZER with 0.20% BARRICADE® herbicide controls certain grass and broadleaf weeds *before* they emerge and provides residual control in established turfgrasses, and ornamentals and other non-crop area. AWARD TURF FERTILIZER with 0.20% BARRICADE herbicide inhibits shoot and root growth in susceptible weeds. AWARD TURF FERTILIZER with 0.20% BARRICADE herbicide will not control established weeds. Weed control is most effective when AWARD TURF FERTILIZER with 0.20% BARRICADE herbicide is activated by at least 0.5 inch of rainfall or irrigation before weed seeds germinate and within 14 days following application.

**Weeds Controlled**

Barnyardgrass	Crowfootgrass	Knotweed	Rescuegrass <sup>4</sup>
Bluegrass, Annual	Cupgrass, Woolly	Kochia	Shepherd's Purse
<i>(Poa annua)</i> <sup>1</sup>	Foxtails, Annual	Lambsquarter, Common	Signalgrass, Broadleaf
Carpetweed	Goosegrass <sup>3</sup>	Lovegrass	Speedwell, Persian
Chickweed, Common	Henbit	Panicum (Texas, Fall, Brown top)	Sprangleop
Chickweed, Mouseear	Itchgrass	Pigweed	Spurge, Prostrate
(from seed)	Johnsongrass	Pigweed	Witchgrass
Crabgrass	(from seed)	Purslane, Common	Woodsoorrel, Yellow
(Large, Smooth) <sup>2</sup>	Junglerice	Pusley, Florida	(from seed)

<sup>1</sup>In areas where *Poa annua* is a winter annual, apply AWARD TURF FERTILIZER with 0.20% BARRICADE herbicide (see rate table) in August or September to established, non-overseeded turf before *Poa annua* seeds germinate. These timings are approximate. Consult State Extension Service for more specific timing for your area.

<sup>2</sup>Fall Applications for Spring Crabgrass Control in Cool-Season Grasses: In those areas where the ground freezes in the winter, AWARD TURF FERTILIZER with 0.20% BARRICADE herbicide can be applied in the fall after the soil temperature falls below 50°F but before the ground freezes. This application will control crabgrass the following spring.

<sup>3</sup>In many areas, a single application will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, make a second application 60-90 days later to ornamentals and lawns (if recommended for your lawn grass species).

<sup>4</sup>Suppression only.

**ESTABLISHED TURF**

**Golf courses (except putting greens), lawns, parks, athletic fields, sod farms How to Apply**


- Do not apply more than 17.2 pounds product/1,000 sq ft (1.5 lb ai/A) per calendar year.
- Do not make more than two applications per calendar year.
- Refer to *Rates for Grass Species* in table and *Granular Spreader Settings* table for spreader settings to deliver the rate recommended for your lawn grass species. Uniform coverage is necessary to provide the most effective level of control.
- Apply AWARD TURF FERTILIZER with 0.20% BARRICADE herbicide to the turf in the early spring prior to crabgrass emergence. See *Crabgrass Germination Dates* in the map below for the approximate timing in your area. Always apply before weeds emerge. AWARD TURF FERTILIZER with 0.20% BARRICADE herbicide will not control emerged crabgrass.
- If you have a known serious weed problem or live in an area with an extended growing season, you may need to make a second application 60-90 days later.

• If you live in Zones D and E (see *Crabgrass Germination Dates* map), you may apply a higher rate (Medium or High Rate) or make two applications for more effective crabgrass control, if recommended for your turfgrass species.

• Do not make a second application to Fine Fescue, Creeping Bentgrass, and newly sprigged and plugged Bermudagrass. Do not make a second application if you applied the High Rate to Buffalograss, Kentucky Bluegrass, or Perennial Ryegrass.

• For best control - After application, water in with ½ inch of water within 14 days, either from watering or rain.

**Crabgrass Seed Germination Dates**



Rates for Lawn Grass Species	Application Schedule <sup>1</sup>	Application Rate <sup>2</sup>		
		Base	Medium	High
Bermudagrass (established) Bahagrass Centipedegrass Kikuyugrass Seashore Paspalum St. Augustine grass Tall Fescue (including turf-type) Zoysiagrass	First Application	Base	Medium	High
	Second Application (optional)	Base or Medium or High	Base or Medium or High	Base or Medium or High
Buffalograss Kentucky Bluegrass Perennial Ryegrass	First Application	Base	Medium	High
	Second Application (optional)	Base or Medium	Base or Medium	—
Fine Fescue (including Red Fescue)	One Application Only	Base or Medium or High	—	—
Creeping Bentgrass (0.5 inch and greater height) <sup>3</sup>	One Application Only	Base or Medium	—	—
Bermudagrass (newly sprigged and plugged) <sup>4</sup>	One Application Only	Base or Medium	—	—

<sup>1</sup> A second application may be necessary because of heavy weed pressure or extended growing season.

<sup>2</sup> See *Granular Spreader Settings* table for spreader settings to deliver the rate.

<sup>3</sup> To avoid grass injury, do not apply AWARD TURF FERTILIZER with 0.20% BARRICADE herbicide to creeping bentgrass mowed at less than 0.5 inch in height.

<sup>4</sup> Newly sprigged or plugged Bermudagrass rooting may be temporarily slowed.

Make and Models	0.38 lb ai/A	0.5 lb ai/A	0.75 lb ai/A
	4.4 lb product/1,000 sq ft Base Rate	5.7 lb product/1,000 sq ft Medium Rate	8.6 lb product/1,000 sq ft High Rate
<b>Spyker 24</b>	4.5	4.75	5.5
<b>Scotts R8A</b>	L	N	P
<b>Lesco Rotary</b>	J-K	L	N
<b>Prizelawn CB Rotary</b>	5	5.25	6
<b>Earthway 2400 Rotary</b>	17	18	20
<b>Lely Broadcast</b>	3.5	3.8	4.2

<sup>1</sup> Assume 3 mph walking speed.

<sup>2</sup> It is recommended that you calibrate your spreader before application since manufacturer's settings can vary among the same model of spreader.

<sup>3</sup> Do not apply more than 1.5 lb ai/A per year.

**Net Wt. 50 lbs. (22.7 kg.)**

### Use Precautions

- Do not apply to putting greens or turfgrass areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa annua*) are desirable species.
- Do not cut (harvest) treated sod before 90 days after application. To avoid turfgrass injury, do not apply to newly-set sod until the sod has rooted into the second inch of soil and exposed edges have filled in.
- Application of AWARD TURF FERTILIZER with 0.20% BARRICADE herbicide to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.
- **Seeding:** Do not apply to overseeded areas for 60 days after seeding, or until after the second mowing, whichever is longer. Do not overseed for 4 months after last applications of this product.

### LANDSCAPE, FIELD-GROWN ORNAMENTALS AND NON-CROP AREAS

#### How to Apply

- Apply AWARD TURF FERTILIZER with 0.20% BARRICADE herbicide as a broadcast, over-the-top application at 7.5-17.2 lb product/1,000 sq ft. Use higher rates when weed pressure was a severe problem in the past.
- Make a single application in fall and/or spring prior to weed emergence. Another application may be made 60-90 days later, if necessary. Do not exceed a total of 17.2 lb product/1,000 sq ft per year.
- After application, water in with ½ inch of water within 14 days, either from watering or rain, or incorporate into the top 1-2 inches of soil within 14 days after application.
- On established ornamentals, watering or rainfall soon after application will wash residues off plant foliage and activate AWARD TURF FERTILIZER with 0.20% BARRICADE herbicide in the soil.
- On newly-transplanted ornamentals, apply after the soil has settled around the transplants.
- For best weed control, apply to soil free of clods, weeds, and debris such as leaves.
- Prior to application, control existing weeds by hand weeding, cultivation, or by the use of an herbicide for emerged weeds.
- Plants under stress (such as heat, drought, or frost damage) may be sensitive to AWARD TURF FERTILIZER with 0.20% BARRICADE herbicide.

### Tolerant Ornamentals

AWARD TURF FERTILIZER with 0.20% BARRICADE herbicide will not harm most trees, shrubs, vines and flowers.

Table 1 - Tolerant Ornamentals - All States

Scientific name	Common name
<i>Abies</i> spp.	Fir species (Balsam, Fraser, Noble, etc.)
<i>Acer palmatum</i>	Japanese Maple
<i>Acer platanoides</i>	Norway Maple
<i>Actinidia chinensis</i>	Kiwi*
<i>Agapanthus africanus</i>	Lily-of-the-Nile (African Lily)
<i>Arctostaphylos densiflora</i>	Vine Hill Manzanita
<i>Arctotheca calendula</i>	Cape Weed
<i>Aucuba japonica</i>	Japanese Aucuba
<i>Berberis gladiolifolia</i>	Barberry
<i>Berberis julianae</i>	Wintergreen Barberry
<i>Berberis mentorensis</i>	Mentor Barberry
<i>Berberis thunbergii</i>	Japanese Barberry
<i>Berberis verruculosa</i>	Warty Barberry
<i>Buxus microphylla</i>	Japanese Boxwood
<i>Callistemon viminalis</i>	Weeping Bottlebrush
<i>Calluna vulgaris</i>	Scotch Heather
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)
<i>Cassia artemisioides</i>	Feathery Cassia
<i>Ceanothus rigidus</i>	Wild Lilac
<i>Chamaecyparis pisifera</i>	False Cypress
<i>Cleyera japonica</i>	Cleyera
<i>Citrus</i> spp.	Citrus species*
<i>Cornus florida</i>	Flowering Dogwood
<i>Cornus stolonifera</i>	American Dogwood
<i>Cortaderia selloana</i>	Pampas Grass
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster
<i>Cotoneaster buxifolius</i>	Cotoneaster
<i>Cotoneaster dammeri</i>	Bearberry Cotoneaster
<i>Cotoneaster microphyllus</i>	Rockspray Cotoneaster
<i>Crataegus</i> spp.	Hawthorne
<i>Cupressus sempervirens</i>	Italian Cypress

### Scientific name

*Delosperma alba*  
*Dodonaea viscosa*  
*Elaeagnus pungens*  
*Euonymus fortunei*  
*Euonymus japonica*  
*Euonymus kiautschovicks*  
*Fatsia japonica*  
*Forsythia intermedia*  
*Forsythia viridissima*  
*Gardenia jasminoides*  
*Gladiolus* spp.  
*Hedera helix*  
*Hibiscus*  
*Hibiscus Rosa-sinensis*  
*Ilex cornuta*  
*Ilex crenata*  
*Ilex opaca*  
*Ilex pernyi*  
*Ilex vomitoria*  
*Iris* spp.  
*Jasminium nudiflorum*  
*Juniperus chinensis*  
*Juniperus conferta*  
*Juniperus horizontalis*  
*Juglans* spp.  
*Justicia brandegeana*  
*Lagerstromia indica*  
*Ligustrum amurense*  
*Ligustrum japonicum*  
*Ligustrum lucidum*  
*Liriope muscari*  
*Lonicera japonica*  
*Lonicera tatarica*  
*Magnolia* spp.  
*Malephora luteola*  
*Malus* spp.  
*Nandina domestica*  
*Narcissus* spp.  
*Nerium* spp.  
*Olea europaea*  
*Ophiopogon japonicus*  
*Osteospermum fruticosum*  
*Oxydendrum arboreum*  
*Persea americana*  
*Photinia fraseri*  
*Picea* spp.  
*Pieris japonica*  
*Pinus brutia*  
*Pinus canariensis*  
*Pinus elliotii*  
*Pinus halepensis*  
*Pinus nigra*  
*Pinus palustris*  
*Pinus radiata*  
*Pinus strobus*  
*Pinus sylvestris*  
*Pinus taeda*  
*Pinus thunbergiana*  
*Pinus virginiana*  
*Pistacia* spp.  
*Pittosporum thombifolium*  
*Pittosporum tobira*  
*Podocarpus macrophyllus*  
*Prunus laurocerasus*  
*Prunus* spp.

*Pseudotsuga menziesii*  
*Pyracantha coccinea*  
*Pyracantha fortuneana*  
*Pyracantha koidzumii*

### Common name

White Trailing Ice Plant  
Hop Bush  
Silverberry  
Wintercreeper  
Japanese Spindle Tree (Evergreen Euonymus)  
Spreading Euonymus  
Japanese Aralia  
Border Forsythia  
Greenstem Forsythia  
Gardenia, Cape-Jasmine  
*Gladiolus* species  
English Ivy  
Rose of Sharon  
Chinese Hibiscus  
Chinese Holly  
Japanese Holly  
American Holly  
Holly  
Yaupon Holly  
*Iris* species  
Winter Jasmine  
Chinese Juniper  
Shore Juniper  
Creeping Juniper  
Walnut\*  
Shrimp Plant  
Crape Myrtle  
Amur Privet  
Japanese Privet  
Glossy Privet (Wax-Leaf)  
Big Blue Lilyturf  
Japanese Honeysuckle  
Tatarian Honeysuckle  
Magnolia species  
Ice Plant  
Crabapple\*  
Heavenly Bamboo  
Narcissus species  
Oleander  
Olive\*  
Mondo Grass  
Trailing African Daisy  
Sourwood  
Avocado\*  
Frasier's Photinia (Redtip)  
Spruce species (Colorado Blue, Norway, etc.)  
Lily-of-the-Valley Shrub  
Calabrian Pine  
Canary Island Pine  
Slash Pine  
Aleppo Pine  
Austrian Black Pine  
Longleaf Pine  
Monterey Pine  
Eastern White Pine  
Scotch Pine  
Loblolly Pine  
Japanese Black Pine  
Virginia Pine  
Pistachio\*  
Queensland Pittosporum  
Japanese Pittosporum  
Japanese Yew  
English Laurel  
Almond, Apricot, Nectarine, Peach, Plum and Prune\*  
Douglas Fir  
Firethorn Scarlet  
Firethorn  
Firethorn

### Scientific name

*Pyrus* spp.  
*Quercus rubra*  
*Rhaphiolepis indica*  
*Rhododendron (including Azalea)*

*Rosa banksiae*  
*Rosmarinus officinalis*  
*Rumohra adiantiformis*  
*Santolina virens*  
*Sedum album*  
*Syzygium paniculatum*  
*Taxus cuspidata*  
*Taxus media*  
*Thuja occidentalis*  
*Trachelospermum asiaticum*  
*Tsuga canadensis*  
*Tulipa* spp.  
*Viburnum japonicum*  
*Viburnum odoratissimum*  
*Viburnum plicatum*  
*Viburnum rigidum*  
*Viburnum tinus*  
*Viburnum trilobum*  
*Viburnum wrightii*  
*Vinca major*  
*Vinca minor*  
*Vitis* spp.  
*Weigela florida*  
*Yucca aloifolia*  
*Yucca filamentosa*

### Common name

Bradford Pear spp.  
Oak species  
Indian Hawthorne  
'Coral Bells'  
'Formosa'  
'Hino-crimson'  
'PJM'  
'Roseum Elegans'  
Lady Bank's Rose  
Rosemary\*  
Leatherleaf Fern

Stonecrop  
Japanese Boxcherry  
Japanese Yew  
Yew  
American Arborvitae  
Star Jasmine  
Canada Hemlock  
Tulip species  
Japanese Viburnum  
Sweet Viburnum  
Japanese Snowball  
Canary Island Viburnum  
Laurustinus  
Cranberry Bush  
Leatherleaf Viburnum  
Vinca  
Dwarf Periwinkle  
Grape\*  
Old Fashioned Weigela  
Spanish Bayonet  
Yucca, Adam's Needle

\*Do not use on food producing trees, vines, or plants.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

### Pesticide Storage

Store in a cool, dry place. Keep container closed and away from food, feedstuffs, and domestic water supplies.

### Pesticide Disposal

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

### Container Handling

Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Then offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### Pesticide Disposal and Container Handling [Household/Residential]

Nonrefillable container. Do not reuse or refill this container.

**If empty:** Place in trash or offer for recycling if available.

**If partly filled:** Call your local solid waste agency for disposal instructions.

Never place unused product down any indoor or outdoor drain.

Award® and Barricade® are registered trademarks of a Syngenta Group Company.

# EC GROW

Distributed and Guaranteed By Eau Claire Co-op Oil Co.  
P.O. Box 837, Eau Claire, WI 54702

Doc.# RP100-015-15